

REPRESENTATIONS OF NON-COMMUTATIVE QUANTUM GROUPS

BENOIT KRIEGK AND MICHEL VAN DEN BERGH

ABSTRACT. We discuss the representation theory of the bialgebra $\underline{\text{end}}(A)$ introduced by Manin. As a side result we give a new proof that Koszul algebras are distributive and furthermore we show that some well-known N -Koszul algebras are also distributive.

1. INTRODUCTION

Throughout we work over a ground field k . If A is a $\mathbb{Z}$ -graded algebra then there is a bialgebra $\underline{\text{end}}(A)$ coacting on A which is universal in an appropriate sense (see [14] and also §3 below). For example if $A = k[x, y]$ then

$$(1.1) \quad \underline{\text{end}}(A) = k[a, b, c, d]/(ac - ca, ad - da + bc - cb, bd - db)$$

and the comultiplication and coaction can be written concisely as

$$\begin{aligned} \Delta \begin{pmatrix} a & b \\ c & d \end{pmatrix} &= \begin{pmatrix} a & b \\ c & d \end{pmatrix} \otimes \begin{pmatrix} a & b \\ c & d \end{pmatrix} \\ \delta \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} a & b \\ c & d \end{pmatrix} \otimes \begin{pmatrix} x \\ y \end{pmatrix} \end{aligned}$$

We see in particular that there is a bialgebra morphism from $\underline{\text{end}}(A)$ to $\mathcal{O}(M_{2 \times 2})$, the coordinate ring of the monoid of 2×2 -matrices. So is tempting in this case to think of the bialgebra $\underline{\text{end}}(A)$ as a non-commutative version of this monoid.

Recently the bialgebra $\underline{\text{end}}(A)$ appeared as a vehicle for deriving combinatorial identities such as the quantum MacMahon Master theorem [12, 13]. In order to understand how far such methods can be pushed it is useful to describe the representation theory of $\underline{\text{end}}(A)$.¹ In this paper we will accomplish this for a large class of algebras.

Our results will be for so-called *distributive algebras*. Let $A = TV/(R)$ with $R \subset V^{\otimes N}$. We say that A is distributive if for all n the subspaces $V^{\otimes a} \otimes R \otimes V^{\otimes n-N-a}$ of $V^{\otimes n}$ generated a distributive lattice. If $N = 2$ then it is well known that distributivity is the same as Koszulity [4] (in Theorem 2.1 below we give an alternative proof of this result). For $N > 2$ distributivity is related to “ N -Koszulity”, a higher Koszul property introduced by Berger [6].

Recall that A is N -Koszul if the free modules occurring in the minimal resolution of k are generated in degrees $0, 1, N-1, N, 2N-1, 2N$, etc. . . . One has the following

1991 *Mathematics Subject Classification.* Primary 16T10; Secondary 16S37.

Key words and phrases. Quantum groups, N -Koszulity, distributivity.

The second author is a senior researcher at the FWO.

¹Unless otherwise specified, representations of a bialgebra are coalgebra representations.

implications

$$(1.2) \quad \text{Distributivity} + \text{“extra condition”} \Rightarrow N\text{-Koszul} \Rightarrow \text{“extra condition”}$$

Here the “extra condition” is a supplementary condition which is vacuous for $N = 2$ and for $N = 3$ reduces to

$$R \otimes V \otimes V \cap V \otimes V \otimes R \subset V \otimes R \otimes V$$

It is unfortunately unknown if the first implication in (1.2) is reversible.

To state our results we recall the definition of the category Cube_n introduced by Polishchuk and Positselski in [15]. Cube_0 is the category of vector spaces, Cube_1 is the category of vector space homomorphisms, Cube_2 is the category of commutative squares etc. . . . In general Cube_n is given by the representations of a hypercube Q_n with commutative faces. We put $\text{Cube}_\bullet = \bigoplus_n \text{Cube}_n$. Then as observed in [15] $\text{Cube}_\bullet$ carries a natural monoidal structure. See §2.1.

The following is our main result concerning the representation theory of $\underline{\text{end}}(A)$.

Proposition 1.1. *(Proposition 4.3 below) Assume that A is distributive. Then there is a monoidal functor $\mathcal{F}_A : \text{Cube}_\bullet \rightarrow \text{CoMod}(\underline{\text{end}}(A))$ which induces an equivalence of monoidal categories*

$$\bar{\mathcal{F}}_A : \text{Cube}_\bullet / \mathcal{S}_A \cong \text{CoMod}(\underline{\text{end}}(A))$$

where $\mathcal{S}_A$ is the localizing subcategory of $\text{Cube}_\bullet$ generated by the simples S such that $\bar{\mathcal{F}}_A(S) = 0$.

This proposition yields a description of the representation theory of $\underline{\text{end}}(A)$ in terms of the subquivers of $(Q_n)_n$ whose vertices correspond to the simples that are not annihilated by $\mathcal{F}_A$.

It remains to determine the simples that are killed by $\mathcal{F}_A$. If we assume the extra condition then this is a tractable problem.

Proposition 1.2. *(Corollary 4.4 below) Assume that A satisfies the extra condition (besides being distributive). Then the monoidal category $\text{CoMod}(\underline{\text{end}}(A))$ is entirely determined by the dimensions of the vector spaces*


$$\bigcap_a V^{\otimes a} \otimes R \otimes V^{\otimes n-a-N}$$

If $N = 2$ then $\text{CoMod}(\underline{\text{end}}(A))$ is entirely determined by the Hilbert series of A .

In §5 we discuss the case of the symmetric algebra. We prove

Proposition 1.3. *(Proposition 5.1 below) Let $A = SV$ where V is an n -dimensional vector space. Then the simple $\underline{\text{end}}(A)$ -representations when viewed as $M_{n \times n}$ representations are of the form $S^{\lambda/\mu}V$ with λ/μ running through the set of rim hooks.*

Recall that a rim hook is a connected skew diagram containing no 2×2 -squares. An example is the following



(1.3)

In the last section of this paper we discuss some examples of N -Koszul algebras which we can show to be distributive. It is noteworthy that we do not know an example of an N -Koszul algebra which is not distributive. On the other hand we also do not know (for $N > 2$) a homological characterization of distributivity.

We show distributivity of the following algebras

- Algebras derived from Koszul algebras. This is more or less a tautology but it gives a quick proof that the generalized symmetric algebra $TV/\wedge^N V$ introduced by Berger is N -Koszul.
- The deformed Yang-Mills algebras introduced by Connes and Dubois-Violette in [9, 10]. We show that these algebras are in fact “confluent” (see [5] and also §6.2 below).
- The 3-dimensional cubic Artin-Schelter regular algebras of “Type A”. This proof also relies on confluence but in a more sophisticated way. We rely on a generalization of the I -type condition which was introduced in [18].

2. THE CATEGORY Cube_n

2.1. Generalities. The quiver Q_n is the quiver with relations which has a vertex x_I for every $I \subset \{1, \dots, n\}$ and arrows $x_{IJ} : x_I \rightarrow x_J$ for every $I \subset J$. The relations are given by $x_{JK}x_{IJ} = x_{IK}$ (we write paths in functional notation). If $n \leq 0$ then we declare Q_n to be a single vertex $x_\emptyset$ without arrows.

We denote the projective Q_n representation corresponding to x_I by $P_I^{(n)}$ and the corresponding simple representation by $S_I^{(n)}$. The category of Q_n -representations (not necessarily finite dimensional) is denoted by Cube_n . The full subcategory consisting of finite dimensional representations is denoted by cube_n . An object in Cube_n is generally written as $(X_I)_I$ where I runs through the subsets of $\{1, \dots, n\}$. We recall the following result.

Theorem 2.1.1. [15, Lemma 9.1] *A collection of subobjects $R_1, \dots, R_n \subset X$ of an object X in a k -linear abelian category $\mathcal{C}$ generates a distributive lattice iff the right exact functor $F : \text{cube}_n \rightarrow \mathcal{C}$ defined by $F(P_\emptyset) = X$ and $F(P_{\{i\}} \rightarrow P_\emptyset) = R_i \hookrightarrow X$ is exact.*

Note that if $\mathcal{C}$ is a Grothendieck category then the functor F extends to a right exact functor $F : \text{Cube}_n \rightarrow \mathcal{C}$ commuting with filtered exact limits. If $F : \text{cube}_n \rightarrow \mathcal{C}$ is exact then so is its extension.

As observed in [15] the categories Cube_n carry an interesting monoidal structure. We first consider the bifunctor $(m, n \geq 0)$

$$\boxtimes : \text{Cube}_m \times \text{Cube}_n \rightarrow \text{Cube}_{m+n} : ((X_I)_I, (Y_J)_J) \mapsto (X_I \otimes Y_J)_{I \cup J + m}$$

where $J + m \subset \{m+1, \dots, m+n\}$ is the translation of J by m . This tensor product is clearly associative and biexact. In addition we have $P_I^{(m)} \boxtimes P_J^{(n)} = P_{I \cup J + m}^{(m+n)}$. For a number $N \geq 2$ (which will remain constant) one also defines a shifted tensor product

$$\boxtimes : \text{Cube}_m \times \text{Cube}_n \rightarrow \text{Cube}_{m+n+N-1} : ((X_I)_I, (Y_J)_J) \mapsto (X_I)_I \boxtimes P_\emptyset^{(N-1)} \boxtimes (Y_J)_J$$

This tensor product is still associative. On the level of projectives we have

$$(2.1) \quad P_I^{(m)} \otimes P_J^{(n)} = P_{I \cup (J+(m+N-1))}^{(m+n+N-1)}$$

We extend this formula beyond its original context by declaring it to be valid for $m \geq -N + 1$, $n \geq -N + 1$ where we recall that by our conventions if $m \leq 0$ then $P_I^{(m)}$ exists only for $I = \emptyset$. Concretely we have for $m = -N + 1, \dots, 0$, $n \geq 0$

$$\otimes : \text{Cube}_m \times \text{Cube}_n \rightarrow \text{Cube}_{m+n+N-1} : (P_\emptyset^{(m)}, (Y_J)_J) \mapsto P_\emptyset^{(m+N-1)} \boxtimes (Y_J)_J$$

for $m \geq 0$, $n = -N + 1, \dots, 0$:

$$\otimes : \text{Cube}_m \times \text{Cube}_n \rightarrow \text{Cube}_{m+n+N-1} : ((X_I)_I, P_\emptyset^{(n)}) \mapsto (X_I)_I \boxtimes P_\emptyset^{(n+N-1)}$$

and for $m = -N + 1, \dots, 0$, $n = -N + 1, \dots, 0$:

$$\otimes : \text{Cube}_m \times \text{Cube}_n \rightarrow \text{Cube}_{m+n+N-1} : (P_\emptyset^{(m)}, P_\emptyset^{(n)}) \mapsto P_\emptyset^{(m+n+N-1)}$$

If $(\mathcal{A}_i)_i$ is a family of abelian categories then we denote by $\bigoplus_i \mathcal{A}_i$ the abelian category whose objects are formal direct sums $\bigoplus_i A_i$ with $A_i \in \mathcal{A}_i$. The Hom-sets are given by

$$\text{Hom}_{\mathcal{A}} \left(\bigoplus_i A_i, \bigoplus_i B_i \right) = \prod_i \text{Hom}_{\mathcal{A}_i}(A_i, B_i)$$

By $\bigoplus'_i \mathcal{A}_i$ we mean the full subcategory of $\bigoplus_i \mathcal{A}_i$ where we only consider objects $(A_i)_i$ with a finite number of non-zero A_i .

Put $\text{Cube}_\bullet = \bigoplus_{n \geq -N+1} \text{Cube}_n$ and $\text{cube}_\bullet = \bigoplus'_{n \geq -N+1} \text{cube}_n$. With the tensor product defined above $\text{Cube}_\bullet$ and $\text{cube}_\bullet$ are monoidal categories with a biexact tensor product. The unit object is $P_\emptyset^{(-N+1)}$.

2.2. Tensor products of simples. For the benefit of the reader we discuss the tensor product of simples in the category $\text{Cube}_\bullet$.

Proposition 2.2.1. *Let $I \subset \{1, \dots, m\}$, $J \subset \{1, \dots, n\}$ with the convention that if $m \leq 0$ or $n \leq 0$ then correspondingly $I = \emptyset$ or $J = \emptyset$. Put $J' = J + m + N - 1$. The simples occuring in the Jordan-Holder series of $S_I^{(m)} \otimes S_J^{(n)}$ are $S_K^{(m+n+N-1)}$ with $K = I \cup J' \cup C$ for $C \subset \{\max(1, m+1), \dots, \min(m+N-1, m+n+N-1)\}$. The simples occur with multiplicity one and are ordered according to $|K|$ with $S_{I \cup J'}^{(m+n+N-1)}$ appearing on top.*

Proof. We will consider the case $m, n \geq 1$. The other cases are similar. We recall the formula for the tensor product. Let $X \in \text{Cube}_m$, $Y \in \text{Cube}_n$. Then

$$X \otimes Y = X \boxtimes P_\emptyset^{(N-1)} \boxtimes Y$$

If we apply this with $X = S_I^{(m)}$, $Y = S_J^{(n)}$ we find for $K \subset \{1, \dots, m\}$

$$(S_I^{(m)} \otimes S_J^{(n)})_K = \begin{cases} k & \text{if } K = I \cup J' \cup C \text{ with } C \subset \{m+1, \dots, m+N-1\} \\ 0 & \text{otherwise} \end{cases}$$

So the simples that occur in the Jordan-Holder series of $S_I^{(m)} \otimes S_J^{(n)}$ are indeed as indicated. $\square$

2.3. Koszulity and Cube $_{\bullet}$. It was observed in [15] that when $N = 2$ the category $\text{Cube}_{\bullet}$ is intimately connected with the Koszul property for graded algebras. Similarly, as we show in this section, when $N > 2$, the category $\text{Cube}_{\bullet}$ is connected with the N -Koszul property introduced by Roland Berger in [6].

Let $N \geq 2$ and consider $A = T(V)/(R)$ with V a finite dimensional k -vectorspace and $R \subseteq V^{\otimes N}$. We put

$$R_i^{(n)} = V^{\otimes i-1} \otimes R \otimes V^{\otimes n-N-i+1} \subset V^{\otimes n}$$

and for $I \subset \{1, \dots, n - N + 1\}$

$$R_I^{(n)} = \bigcap_{i \in I} R_i^{(n)}$$

We define F_A as the right exact functor commuting with filtered colimits

$$F_A : \text{Cube}_{n-N+1} \rightarrow \text{Vect}$$

which sends $P_I^{(n-N+1)}$ to $R_I^{(n)}$. As usual we denote its left derived functor by LF_A . It is easy to see that F_A is a monoidal functor.

Let $\mathcal{L}_n(A)$ be the lattice of subspaces of $V^{\otimes n}$ generated by $(R_i^{(n)})_i$.

Definition 2.3.1. A is *distributive* if for each n the lattice $\mathcal{L}_n(A)$ is distributive.

Definition 2.3.2. [6] A satisfies the *extra condition* if for all $1 \leq i \leq l \leq j \leq n$ with $j \leq i + N - 1$ we have

$$R_i^{(n)} \cap R_j^{(n)} \subset R_l^{(n)}$$

For use below we also put for $n \geq N$

$$J_n = R_1^{(n)} \cap \dots \cap R_{n-N+1}^{(n)}$$

Thus $J_N = R$. For $n \leq N - 1$ we put $J_n = V^{\otimes n}$. We denote the composition

$$(2.2) \quad A \otimes J_n \rightarrow A \otimes V \otimes J_{n-1} \rightarrow A \otimes J_{n-1}$$

by δ . Here the first map is derived from the inclusion $J_n \subset V \otimes J_{n-1}$ and the second map is derived from the multiplication $A \otimes V \rightarrow A$. Note that $\delta^N = 0$. Thus $(A \otimes J_n, \delta)$ is an N -complex [7].

Definition 2.3.3. [6] A is N -Koszul if the following complex

$$(2.3) \quad \dots \rightarrow A \otimes J_{2N} \xrightarrow{\delta^{N-1}} A \otimes J_{N+1} \xrightarrow{\delta} A \otimes R \xrightarrow{\delta^{N-1}} A \otimes V \xrightarrow{\delta} A \rightarrow k$$

is exact.

A famous theorem by Backelin [4] states that if $N = 2$ then A is distributive if and only if it is Koszul (see below for a proof in the current setting). In general Berger proved the implications [6]

$$(2.4) \quad \text{Distributivity} + \text{extra condition} \Rightarrow N\text{-Koszul} \Rightarrow \text{extra condition}$$

(the extra condition is vacuous if $N = 2$). It is unknown if the converse of the first implication holds.

The category $\text{Cube}_{\bullet}$ contains a preimage under F_A of the degree m part of the Koszul complex of A (2.3). Put

$$W_{m,n} = S_{\emptyset}^{(m-N+1)} \otimes P_{\{1, \dots, n-N+1\}}^{(n-N+1)}$$

Since $A_m = F_A(S_\emptyset^{(m-N+1)})$ and $J_n = F_A(P_{\{1, \dots, n-N+1\}}^{(n-N+1)})$ (with the usual caveat if $n - N + 1 \leq 0$) one obtains

$$F_A(W_{m,n}) = A_m \otimes J_n$$

A quick computation shows

$$(2.5) \quad (W_{m,n})_I = \begin{cases} k & \text{if } \{m+1, \dots, m+n-N+1\} \subset I \subset \{m-N+2, \dots, m+n-N+1\} \\ 0 & \text{otherwise} \end{cases}$$

and all maps are either the identity (when they are between copies of k) or zero. Define

$$\delta : W_{m,n} \rightarrow W_{m+1,n-1}$$

such that δ_I is the identity on k when $(W_{m,n})_I = (W_{m-1,n+1})_I = k$ and zero otherwise. One checks that this is indeed a well defined map in $\text{Cube}_\bullet$ and that $F_A(\delta) = \delta_m$ where δ_m is the degree m part of the map (2.2). Furthermore $\delta^N = 0$. It follows that the following complex in $\text{Cube}_\bullet$

$$(2.6) \quad \cdots \xrightarrow{\delta} W_{m-2N,2N} \xrightarrow{\delta^{N-1}} W_{m-N-1,N+1} \xrightarrow{\delta} W_{m-N,N} \xrightarrow{\delta^{N-1}} W_{m-1,1} \xrightarrow{\delta} W_{m,0} \rightarrow S_\emptyset^{(-N+1)} \cdot \delta_{m,0}$$

maps to the degree m part of the Koszul complex.

Let us say that $I \subset \{1, \dots, m\}$ has a hole of size u (with $u \geq 1$) if there exist $i < j \in I$ such that $j - i = u + 1$ and for all $i < l < j$ we have $l \notin I$.

Proposition 2.3.4. *The homology of (2.6) is an extension of simples $S_K^{(m-N+1)}$ such that K contains holes of size $\leq N - 2$.*

Proof. to prove this we will more generally consider the homology of the complexes

$$(2.7) \quad W_{m,n} \xrightarrow{\delta^a} W_{m+a,n-a} \xrightarrow{\delta^{N-a}} W_{m+N,n-N}$$

for any $0 < a < N$. (2.6) is built up from such complexes for $a = 1, N - 1$.

Let H be the homology of (2.7). From (2.5) we deduce that H_I is non-zero iff the following conditions hold

$$(2.8) \quad \begin{aligned} I &\not\subset \{m+2, \dots, m+n-N+1\} \\ \{m+1, \dots, m+n-N+1\} &\not\subset I \\ \{m+a+1, \dots, m+n-N+1\} &\subset I \subset \{m+a-N+2, \dots, m+n-N+1\} \end{aligned}$$

Let us assume that all sets occurring in these conditions are non-empty. This happens when

$$(2.9) \quad N \leq n - a$$

If I satisfies (2.8)(2.9) then we have the following:

$$\begin{aligned} \exists i : m+a-N+2 \leq i \leq m+1 : i \in I \\ \exists j : m+a \geq j \geq m+1 : j \notin I \end{aligned}$$

Consider the triple $\{i, j, m+a+1\}$. Here $i, m+a+1 \in I$ and $j \notin I$. We clearly have $i < j < m+a+1$ and one computes $m+a+1-i \leq N-1$. Hence I contains a hole of size $\leq N-2$.

Looking at the condition (2.9) we see that the homology of (2.6) is an extension of simples $S_K^{(l)}$ such that K contains holes of size $\leq N-2$, except perhaps at $W_{m-1,1}$ and $W_{m,0}$.

To compute the homology at $W_{m-1,1}$ we have to make the substitutions $m \rightarrow m-N$, $n \rightarrow N$, $a \rightarrow N-1$ in (2.7).

Thus the condition (2.8) becomes
(2.10)

$$\begin{aligned} I \not\subset \{m-N+2, \dots, m-N+1\} &= \emptyset \\ \{m-N+1, \dots, m-N+1\} &= \{m-N+1\} \not\subset I \\ \{m, \dots, m-N+1\} &= \emptyset \subset I \subset \{m-N+1, \dots, m-N+1\} = \{m-N+1\} \end{aligned}$$

The only I 's satisfying the last condition are $\emptyset$, $\{m-N+1\}$. $I = \emptyset$ does not satisfy the first condition and $I = \{m-N+1\}$ does not satisfy the second condition. Hence no I satisfies all three. Thus the homology at $W_{m-1,1}$ is zero.

We leave to the reader the easy verification that the homology of (2.6) at $W_{m,0}$ is also zero. $\square$

Remark 2.3.5. A perhaps better way to understand the complex (2.6) is as follows. Write $\mathcal{V} = P_\emptyset^{(-N+2)} \in \text{Cube}_{-N+2}$ and $\mathcal{R} = P_{\{1\}}^{(1)} \in \text{Cube}_1$. Then there is a unique non-zero map (up to scalar) in Cube_1

$$\mathcal{R} \rightarrow \mathcal{V}^{\otimes N}$$

Let $\mathcal{A} = TV/(\mathcal{R})$ be the corresponding graded algebra object in the monoidal abelian category $\text{Cube}_\bullet$. Then $F_A(\mathcal{A}) = A$. The Koszul complex (2.3) can be defined in an arbitrary abelian monoidal category and in particular it can be defined for $\mathcal{A}$. Then (2.6) is just the degree m part of this generalized Koszul complex.

As we have seen the Koszul complex for $\mathcal{A}$ is not exact. Define

$$\text{Cube}_\bullet^{\text{ec}} = \text{Cube}_\bullet / \mathcal{S}^{\text{ec}}$$

where $\mathcal{S}^{\text{ec}}$ is the localizing subcategory generated by the simples $S_K^{(l)}$ such that K has a hole of size $\leq N-2$. It is easy to see that $\text{Cube}_\bullet^{\text{ec}}$ inherits the monoidal structure of $\text{Cube}_\bullet^{\text{ec}}$.

Then the image of the Koszul complex of $\mathcal{A}$ in $\text{Cube}_\bullet^{\text{ec}}$ is exact. Thus $\mathcal{A}$ is an N -Koszul algebra in the monoidal abelian category $\text{Cube}_\bullet^{\text{ec}}$.

The properties introduced in Definitions 2.3.1-2.3.3 translate nicely into properties of the functor F_A . We say that an object $M \in \text{Cube}_m$ is acyclic if $L_i F_A M = 0$ for $i > 0$. Note the following

Lemma 2.3.6. *If $M \in \text{Cube}_a$ and $N \in \text{Cube}_b$ are acyclic then so is $M \otimes N$.*

Proof. Let $P_\bullet \rightarrow M \rightarrow 0$ and $Q_\bullet \rightarrow N \rightarrow 0$ be minimal projective resolutions of M, N . The exactness of the tensor product implies that $P_\bullet \otimes Q_\bullet$ is a minimal projective resolution of $M \otimes N$. Hence $L_\bullet F_A(M \otimes N)$ is computed by $F_A(P_\bullet \otimes Q_\bullet) = F_A(P_\bullet) \otimes F_A(Q_\bullet)$. Since M and N are acyclic, $F_A(P_\bullet)$, $F_A(Q_\bullet)$ are resolutions of $F_A M$, resp. $F_A N$. Hence $F_A(P_\bullet) \otimes F_A(Q_\bullet)$ is a resolution of $F_A M \otimes F_A N$. $\square$

The following is the main result of this section.

Proposition 2.3.7. (1) *A is distributive if and only if the functor F_A is exact for all n , i.e. if and only if all $S_I^{(n-N+1)}$ are acyclic.*

- (2) *A satisfies the extra condition if and only if for all n one has $F_A(S_I^{(n-N+1)}) = 0$ for those I which have a hole of size $\leq N - 2$.*
- (3) *Assume that A satisfies the extra condition. Then for all n and all I that contain a hole of size $\leq N - 2$ one has that $S_I^{(n-N+1)}$ is acyclic.*
- (4) *Assume that A satisfies the extra condition. Then A is N -Koszul if and only if for all n one has that $S_\emptyset^{(n-N+1)}$ is acyclic.*

Proof. (1) This is just a translation of Theorem 2.1.1.

- (2) The simple $S_I^{(n-N+1)}$ has a presentation

$$\bigoplus_{l \notin I} P_{I \cup \{l\}}^{(n-N+1)} \rightarrow P_I^{(n-N+1)} \rightarrow S_I^{(n-N+1)} \rightarrow 0$$

and thus

$$F_A(S_I^{(n-N+1)}) = R_I^{(n)} \Big/ \sum_{l \notin I} R_{I \cup \{l\}}^{(n)}$$

Assume that the extra condition holds and that I has a hole of size $\leq N - 2$, delimited by $i < j$. Pick $i < l < j$. Then $R_I^{(n)} = R_{I \cup \{l\}}^{(n)}$. Hence $F_A(S_I^{(n-N+1)}) = 0$.

Conversely assume $F_A(S_I^{(n-N+1)}) = 0$ for all I containing a hole of size $\leq N - 2$. Then for such an I we have

$$R_I^{(n)} = \sum_{l \notin I} R_{I \cup \{l\}}^{(n)}$$

Iterating this identity we get

$$R_I^{(n)} = \sum_{i=1}^p R_{J_i}^{(n)}$$

where $(J_i)_i$ is the set of all subsets of $\{1, \dots, n\}$ which contain I and which have no holes of size $\leq N - 2$.

Assume now $I = \{i, j\}$ with $i < j \leq i + N - 1$ and take $i < l < j$. Each of the J_i must contain l . Hence $R_{J_i}^{(n)} \subset R_l^{(n)}$ and thus $R_i^{(n)} \cap R_j^{(n)} = R_I^{(n)} \subset R_l^{(n)}$. Hence the extra condition holds.

- (3) Assume that I contains a hole of length $\leq N - 2$, delimited by $i < j$. Pick $i < l < j$. The minimal resolution $K_\bullet$ of $S_I^{(n-N+1)}$ has the form $\bigoplus_{K \supset I} P_K^{(n-N+1)}$. The $P_K^{(n-N+1)}$ come in pairs $P_{K_1}^{(n-N+1)}, P_{K_2}^{(n-N+1)}$ with $l \notin K_1$ and $K_2 = K_1 \cup \{l\}$. Since the extra condition implies $F_A(P_{K_1}^{(n-N+1)}) = F_A(P_{K_2}^{(n-N+1)})$ we easily deduce that $F_A(K_\bullet)$ is acyclic.
- (4) Assume that A is N -Koszul (and hence satisfies the extra condition by (2.4)). We will show by induction that $S_\emptyset^{(m-N+1)}$ is acyclic. For $m = 0$ this is trivial. So assume $m > 0$. Then (2.6) yields a (finite complex)

$$(2.11) \quad \cdots \rightarrow W_{m-N,N} \rightarrow W_{m-1,1} \rightarrow S_\emptyset^{(m-N+1)} \rightarrow 0$$

By induction and Lemma 2.3.6 $W_{m-a,a}$ is acyclic for $a > 0$. By (2.3) the homology of (2.11) is acyclic as well and killed by F_A . Hence $L_i F_A S_\emptyset^{(m-N+1)}$ for $i > 0$ is computed by the homology of.

$$\cdots \rightarrow F_A W_{m-N,N} \rightarrow F_A W_{m-1,1} \rightarrow S_\emptyset^{(m-N+1)} \rightarrow 0$$

This latter complex is precisely the degree m part of the Koszul complex (2.3) which is acyclic by our hypothesis that A is N -Koszul.

Conversely assume that $S_\emptyset^{(m-N+1)}$ is acyclic for all m . Then the terms of (2.6) are acyclic. Since the homology of (2.6) is acyclic and killed by F_A (by (2.3)) it follows that the complex (2.6) becomes exact after applying F_A . The resulting exact sequence is precisely the degree m part of (2.3). Hence combining all m we find that A is N -Koszul. $\square$

Theorem 2.1. [4] *If $N = 2$ then A is Koszul if and only if it is distributive.*

Proof. Assume $N = 2$. If A is distributive then by applying F_A to the complex (2.6) we obtain the degree m part of the Koszul complex of A . Using (1,2,3) of Proposition 2.3.7 we see this complex is exact.

We now prove the “difficult” direction. Assume that A is Koszul. By Proposition 2.3.7 we have to prove that all $S_I^{(n-1)}$ are acyclic. By 2.3.7(4) we know that all $S_\emptyset^{(n-1)}$ are acyclic.

We will use induction on $|I|$ and n to prove that $S_I^{(n-1)}$ is acyclic for all I . If $I = \emptyset$ then there is nothing to prove. So assume $l \in I$. Put $I_1 = I \cap \{1, \dots, l-1\}$ and $I_2 = (I \cap \{l+1, \dots, n-1\}) - l$ (I_1, I_2 can be empty). Then by Proposition 2.2.1 we have an exact sequence (see also [15])

$$(2.12) \quad 0 \rightarrow S_I^{(n-1)} \rightarrow S_{I_1}^{(l-1)} \otimes S_{I_2}^{(n-l-1)} \rightarrow S_{I \setminus \{l\}}^{(n-1)} \rightarrow 0$$

By induction $S_{I_1}^{(l-1)}, S_{I_2}^{(n-l-1)}, S_{I \setminus \{l\}}^{(n-1)}$ are all acyclic. Then (2.12) implies that $S_I^{(n-1)}$ is acyclic as well. $\square$

Remark 2.3.8. The first part of the previous proof works for arbitrary N and yields the first implication in (2.4). In fact for a distributive algebra satisfying the extra condition, the Koszul complex (2.3) is only one of many long exact sequences one may construct.

To be more precise we have already noted that $(A \otimes J_n, \delta)_n$ is an N -complex (see [7]). An N -complex

$$\dots \rightarrow Y^i \xrightarrow{\delta} Y^{i+1} \xrightarrow{\delta} Y^{i+2} \xrightarrow{\delta}$$

can be contracted into a number of genuine complexes

$$\dots \rightarrow Y^i \xrightarrow{\delta^a} Y^{i+a} \xrightarrow{\delta^{N-a}} Y^{i+N} \xrightarrow{\delta^a} Y^{i+N+a} \rightarrow \dots$$

By analyzing the proof of Proposition 2.3.4 (in particular the properties of the complexes (2.7) when (2.9) holds) we find that the contracted complexes of $(A \otimes J_n, \delta)$ are exact, except in their initial degrees. For example if $N = 3$ then the complex

$$\dots \rightarrow A \otimes J_{N+2} \xrightarrow{\delta^2} A \otimes R \xrightarrow{\delta} A \otimes V \otimes V \xrightarrow{\delta^2} A$$

becomes exact starting at the term $A \otimes R$.

On the contrary, the contracted complexes are almost never exact in their initial degrees. This has been observed in [7].

3. THE BIALGEBRA $\underline{\text{end}}(A)$

Let A be a $\mathbb{Z}$ -graded algebra. It is easy to see that there exists an algebra B which is universal for the property that there exists an algebra morphism $\delta : A \rightarrow B \otimes A$ such that $\delta(A_n) \subset B \otimes A_n$ for all n . In other words for any algebra morphism $\partial : A \rightarrow C \otimes A$ which preserves the A -grading there is a unique algebra morphism $\gamma : B \rightarrow C$ such that $\partial = (\gamma \otimes \text{id}) \circ \delta$. From the universality property one immediately obtains a bialgebra structure on B . Following Manin [14] we use the notation $\underline{\text{end}}(A)$ for B .

We consider a special case of this construction. Let $N \geq 2$ and consider $A = T(V)/(R)$ with V a finite dimensional k -vectorspace and $R \subseteq V^{\otimes N}$. Then it is not hard to see that

$$\underline{\text{end}}(A) = T(V^* \otimes V)/(\pi_N(R^\perp \otimes R)),$$

where π_N is the shuffle map

$$\begin{aligned} \pi_N : (V^* \otimes V)^{\otimes N} &\rightarrow V^{*\otimes N} \otimes V^{\otimes N} : \\ f_1 \otimes \cdots \otimes f_N \otimes x_1 \otimes \cdots \otimes x_N &\mapsto f_1 \otimes x_1 \otimes \cdots \otimes f_N \otimes x_N \end{aligned}$$

The co-action of $\underline{\text{end}}(A)$ on A , as well as the bialgebra structure on $\underline{\text{end}}(A)$ can be described concretely in terms of generators. Fix a k -basis $(x_i)_i$ for V and let $(x_i^*)_i$ be its dual basis in V^* . Put $z_i^j = x_j^* \otimes x_i$. Then the co-action of $\underline{\text{end}}(A)$ on A is given in terms of generators by

$$(3.1) \quad \delta(x_i) = \sum_k z_i^k \otimes x_k$$

and the bialgebra structure on $\underline{\text{end}}(A)$ is given by

$$(3.2) \quad \begin{aligned} \Delta(z_i^j) &= \sum_k z_i^k \otimes z_k^j \\ \epsilon(z_i^j) &= \delta_i^j \end{aligned}$$

Below we will be interested in the finite dimensional co-representations of the bialgebra $\underline{\text{end}}(A)$. We first note that $\underline{\text{end}}(A)$ is a graded algebra by putting $\deg z_i^j = 1$. We denote its part of degree n by $\underline{\text{end}}(A)_n$. It follows immediately from (3.2) that $\underline{\text{end}}(A)_n$ is stable under Δ and when equipped with the restriction of ϵ it becomes a coalgebra.

We now recall some facts about sums of coalgebras. Let $(C_i, \Delta_i, \epsilon_i)_i$ be a family of coalgebras. Put $C = \bigoplus_i C_i$. Then C is a coalgebra with comultiplication $\Delta(c) = \bigoplus_i \Delta_i(c_i)$ and counit $\epsilon(c) = \sum_i \epsilon_i(c_i)$ for $c = \bigoplus_i c_i$ with $c_i \in C_i$.

Suppose we are given comodules (W_i, δ_i) for C_i . Then clearly $W = \bigoplus_i W_i$ is a C -comodule with coaction $\delta(w) = \bigoplus_i \delta_i(w_i)$ for $w = \bigoplus_i w_i$, $w_i \in W_i$.

Lemma 3.1. *The functor*

$$\Psi : \bigoplus_i \text{CoMod}(C_i) \rightarrow \text{CoMod}(C) : \bigoplus_i W_i \mapsto \bigoplus_i W_i$$

is an equivalence of categories.

Proof. This is well known. If $W \in \text{CoMod}(C)$ then it decomposes as $(C_i)_i$ -comodules via $W = \bigoplus_i p_i(W)$ where $p_i(w) = \sum_w \epsilon_i(w_{(1)})w_{[2]}$ for $\delta(w) = \sum_w w_{(1)} \otimes w_{[2]}$. $\square$

From this proposition we deduce that in order to understand the representation theory of $\underline{\text{end}}(A)$ it is sufficient to understand the representation theory of $\underline{\text{end}}(A)_n$. Since $\underline{\text{end}}(A)_n$ is finite dimensional we have

$$\text{CoMod}(\underline{\text{end}}(A)_n) \cong \text{Mod}(\underline{\text{end}}(A)_n^{*\circ})$$

Hence we have to describe the algebras $\underline{\text{end}}(A)_n^{*\circ}$. We do this next. For $n < N$ define

$$Z_n(A) = \text{End}_k(V^{\otimes n})$$

and for $n \geq N$

$$Z_n(A) = \left\{ \varphi \in \text{End}_k(V^{\otimes n}) \mid \forall i \in \{1, \dots, n - N + 1\} : \varphi(R_i^{(n)}) \subset R_i^{(n)} \right\}$$

where $R_i^{(n)} = V^{\otimes i-1} \otimes R \otimes V^{\otimes n-N-i+1} \subset V^{\otimes n}$.

Proposition 3.2. *We have as algebras*

$$\underline{\text{end}}(A)_n^{*\circ} \cong Z_n(A)$$

Proof. We consider the most difficult case $n \geq N$. In that case the result follows by the following computation.

$$\begin{aligned} \underline{\text{end}}(A)_n^* &= ((V^* \otimes V)^{\otimes n} / (\pi_N(R^\perp \otimes R))_n)^* \\ &= \left((V^* \otimes V)^{\otimes n} / \sum_{i=0}^{n-N} (V^* \otimes V)^i \otimes \pi_N(R^\perp \otimes R) \otimes (V^* \otimes V)^{n-N-i} \right)^* \\ &\cong \left((V^{*\otimes n} \otimes V^{\otimes n}) / \sum_{i=0}^{n-N} V^{*i} \otimes R^\perp \otimes V^{*n-N-i} \otimes V^i \otimes R \otimes V^{n-N-i} \right)^* \\ &\subset (V^{*\otimes n} \otimes V^{\otimes n})^* \end{aligned}$$

Writing $Y_i = V^{*\otimes i} \otimes R^\perp \otimes V^{*n-N-i} \otimes V^i \otimes R \otimes V^{n-N-i}$ we get

$$\begin{aligned} \underline{\text{end}}(A)_n^* &\cong \left(\sum_{i=0}^{n-N} Y_i \right)^\perp \\ &= \bigcap_{i=0}^{n-N} Y_i^\perp \\ &= \{ \varphi \in (V^{*\otimes n} \otimes V^{\otimes n})^* \mid \forall i : \varphi(Y_i) = 0 \} \\ &= \{ \varphi \in \text{End}_k(V^{\otimes n}) \mid \forall i : \varphi(V^i \otimes R \otimes V^{n-N-i}) \subseteq V^i \otimes R \otimes V^{n-N-i} \} \end{aligned}$$

via the canonical isomorphism $(E^* \otimes E)^* \cong \text{End}_k(E)$ (for finite dimensional E).

Here we have to make the small computation that if $X \subseteq E$ and $\Psi \in (E^* \otimes E)^*$ then $\Psi(X^\perp \otimes X) = 0$ if and only if the corresponding element φ in $\text{End}_k(E)$ satisfies $\varphi(X) \subseteq X$.

We claim that the vector space isomorphism (3.3) is compatible with the given algebra structures on both sides, up to exchanging factors. To verify this we may assume that $R = 0$, i.e. $A = TV$. Then we have to show that the following

isomorphism

$$\begin{aligned}
T(V^* \otimes V)_n^* &\cong ((V^* \otimes V)^{\otimes n})^* \\
&\cong ((V^*)^{\otimes n} \otimes V^{\otimes n})^* \\
&\cong V^{\otimes n} \otimes (V^*)^{\otimes n} \\
&\cong \text{End}(V^{\otimes n})^\circ
\end{aligned}$$

is compatible with the algebra structure. We verify this using the explicit bases introduced above. The comultiplication on $T(V^* \otimes V)_n$ is given by

$$\Delta(z_{i_1}^{j_1} \cdots z_{i_n}^{j_n}) = \sum_{k_1 \cdots k_n} z_{i_1}^{k_1} \cdots z_{i_n}^{k_n} \otimes z_{k_1}^{j_1} \cdots z_{k_n}^{j_n}$$

Hence the multiplication on $T(V^* \otimes V)_n^*$ is given by

$$z_{i_1}^{k_1*} \cdots z_{i_n}^{k_n*} \cdot z_{l_1}^{j_1*} \cdots z_{l_n}^{j_n*} = \delta_{l_1}^{k_1} \cdots \delta_{l_n}^{k_n} z_{i_1}^{j_1*} \cdots z_{i_n}^{j_n*}$$

By definition $z_i^j = x_j^* \otimes x_i$. Then z_i^{j*} when considered as an element of $V \otimes V^*$ is given by $x_j \otimes x_i^*$.

It follows that the multiplication on $V^{\otimes n} \otimes (V^*)^{\otimes n}$ is given by

$$(3.4) \quad (x_{k_1} \cdots x_{k_n} \otimes x_{i_1}^* \cdots x_{i_n}^*) \cdot (x_{j_1} \cdots x_{j_n} \otimes x_{l_1}^* \cdots x_{l_n}^*) = \delta_{l_1}^{k_1} \cdots \delta_{l_n}^{k_n} x_{j_1} \cdots x_{j_n} \otimes x_{i_1}^* \cdots x_{i_n}^*$$

Since $x_{j_1} \cdots x_{j_n} \otimes x_{i_1}^* \cdots x_{i_n}^*$ corresponds to the endomorphism of $V^{\otimes n}$ which sends $x_{i_1} \cdots x_{i_n}$ to $x_{j_1} \cdots x_{j_n}$ and all other basis elements to zero, we see that (3.4) corresponds precisely to the opposite multiplication on $\text{End}(V^{\otimes n})$. $\square$

As $\underline{\text{end}}(A)$ is a graded algebra (see above) the multiplication induces vector space homomorphisms

$$\underline{\text{end}}(A)_m \otimes \underline{\text{end}}(A)_n \rightarrow \underline{\text{end}}(A)_{m+n}$$

which one checks to be coalgebra homomorphisms.

Hence there are dual algebra morphisms

$$(3.5) \quad \mu_{m,n} : \underline{\text{end}}(A)_{m+n}^{*\circ} \rightarrow \underline{\text{end}}(A)_m^{*\circ} \otimes \underline{\text{end}}(A)_n^{*\circ}$$

These are described in the next proposition.

Proposition 3.3. *Under the isomorphism $\underline{\text{end}}(A)_n^{*\circ} \cong Z_n(A)$ given in Proposition 3.2 the algebra morphism (3.5) fits in the following commutative diagram*

$$\begin{array}{ccc}
Z_{m+n}(A) & \longrightarrow & Z_m(A) \otimes Z_n(A) \\
\downarrow & & \downarrow \\
\text{End}(V^{\otimes m+n}) & \xrightarrow{\cong} & \text{End}(V^{\otimes m}) \otimes \text{End}(V^{\otimes n})
\end{array}$$

where the lower map is the obvious algebra isomorphisms

Proof. We have to verify the commutativity of the following diagram

$$\begin{array}{ccc}
((V^* \otimes V)^{\otimes(m+n)})^* & \longrightarrow & ((V^* \otimes V)^{\otimes m})^* \otimes ((V^* \otimes V)^{\otimes n})^* \\
\downarrow & & \downarrow \\
\text{End}(V^{\otimes(m+n)})^\circ & \longrightarrow & \text{End}(V^{\otimes m})^\circ \otimes \text{End}(V^{\otimes n})^\circ
\end{array}$$

This is again an easy verification, e.g. using explicit bases. $\square$

By Lemma 3.1,3.2 we have

$$\text{CoMod}(\underline{\text{end}}(A)) = \bigoplus_n \text{CoMod}(\underline{\text{end}}(A)_n) = \bigoplus_n \text{Mod}(\underline{\text{end}}(A)_n^{*\circ}) = \bigoplus_n \text{Mod}(Z_n(A))$$

Furthermore the induced monoidal structure on the righthand side is as follows: let $W_1 \in \text{Mod}(Z_m(A))$, $W_2 \in \text{Mod}(Z_n(A))$. Then $W_1 \otimes W_2 \in \text{Mod}(Z_{m+n}(A))$ is the pullback under $\mu_{m,n}$ of $W_1 \otimes W_2$, considered as $Z_m(A) \otimes Z_n(A)$ module

We deduce the well-know result (see e.g. [13]).

Proposition 3.4. *Let $\mathcal{L}_n(A)$ be the lattice of subspaces of $V^{\otimes n}$ generated by $(R_i^{(n)})_i$. Then all the objects in $\mathcal{L}_n(A)$ are $Z_n(A)$ -modules and hence $\underline{\text{end}}(A)$ -comodules.*

Proof. The $R_i^{(n)}$ are obviously $Z_n(A)$ representations. Hence so are all objects in $\mathcal{L}_n(A)$. $\square$

4. DISTRIBUTIVE ALGEBRAS

Let $A = TV/(R)$ be as above. Below we assume that A is distributive.

For $I \subset \{1, \dots, n - N + 1\}$ we define

$$C_I^{(n)} = R_I^{(n)} \bigg/ \sum_{J \supsetneq I} R_J^{(n)}$$

We will say that $I \subset \{1, \dots, n - N + 1\}$ is *admissible* if $C_I \neq 0$. Strictly speaking these definitions only make sense if $n \geq N$. For $n < N$ we will put $\{1, \dots, n - N + 1\} = \emptyset$, and hence we only consider $I = \emptyset$, $R_\emptyset = V^{\otimes n}$, $C_\emptyset = V^{\otimes n}$.

Proposition 4.1. (1) *The indecomposable projective $Z_n(A)$ -representations are the $R_I^{(n)}$ for I admissible.*

(2) *The simple $Z_n(A)$ -representations are the $C_I^{(n)}$ for I admissible.*

(3) *$R_I^{(n)}$ is the projective cover of $C_I^{(n)}$.*

(4) *For admissible I, J we have*

$$\text{Hom}_{Z_n(A)}(R_J^{(n)}, R_I^{(n)}) = \begin{cases} k & \text{if } J \supset I \\ 0 & \text{otherwise} \end{cases}$$

When $J \supset I$, the generator of $\text{Hom}_{Z_n(A)}(R_J^{(n)}, R_I^{(n)})$ is given by the inclusion $R_J^{(n)} \subset R_I^{(n)}$.

Proof. Since the $(R_i^{(n)})_i$ generate the distributive lattice $\mathcal{L}_n(A)$ there exists a basis $(w_\alpha)_\alpha$ for $V^{\otimes n}$ such that each vector space in $\mathcal{L}_n(A)$ is spanned by a subset of the w_α [15, Prop. 7.1].

Let e_α be the primitive idempotent in $\text{End}(V^{\otimes n})$ corresponding to the basis vector w_α . I.e. e_α is the projection on kw_α . Then e_α preserves all $R_i^{(n)}$ and hence $e_\alpha \in Z_n(A)$. Then the $(e_\alpha)_\alpha$ form still a maximal set of orthogonal idempotents in $Z_n(A)$.

For $I \subset \{1, \dots, n - N + 1\}$ let $\tilde{C}_I^{(n)}$ be the subspace of $R_I^{(n)}$ spanned by the w_α which are not in some R_J , $J \supsetneq I$. Clearly $C_I^{(n)} \cong \tilde{C}_I^{(n)}$ and hence $\tilde{C}_I^{(n)}$ is non-zero if and only if I is admissible.

We have

$$R_I^{(n)} = \bigoplus_{J \supset I} \tilde{C}_J^{(n)}$$

It follows that

$$\text{End}(V^{\otimes n}) = \bigoplus_{I, J} \text{Hom}(\tilde{C}_I^{(n)}, \tilde{C}_J^{(n)})$$

and

$$(4.1) \quad Z_n(A) = \bigoplus_{J \supset I} \text{Hom}(\tilde{C}_I^{(n)}, \tilde{C}_J^{(n)})$$

Thus

$$Z_n(A) / \text{rad}(Z_n(A)) = \bigoplus_{I \text{ adm.}} \text{Hom}(\tilde{C}_I^{(n)}, \tilde{C}_I^{(n)})$$

For every admissible I pick an α_I such that $w_{\alpha_I} \in \tilde{C}_I^{(n)}$. Then the indecomposable projectives in $Z_n(A)$ -modules are given by

$$Q_I = Z_n(A)e_{\alpha_I}$$

We find

$$Q_I = Z_n(A)e_{\alpha_I} = \bigoplus_{J \supset I} \text{Hom}(kw_{\alpha_I}, \tilde{C}_J^{(n)}) = \text{Hom}(kw_{\alpha_I}, R_I^{(n)}) \cong R_I^{(n)}$$

which proves (1). We also find using the decomposition (4.1)

$$\begin{aligned} \text{Hom}_{Z_n(A)}(Q_J, Q_I) &= e_{\alpha_J} Z_n(A) e_{\alpha_I} \\ &= \begin{cases} k & \text{if } J \supset I \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

which proves (4).

The simple top of Q_I for I admissible may be computed as the cokernel of

$$\bigoplus_{Q_J \not\cong Q_I} Q_J \otimes \text{Hom}_{Z_n(A)}(Q_J, Q_I) \rightarrow Q_I$$

which is the same as the cokernel of

$$\bigoplus_{R_J^{(n)} \subsetneq R_I^{(n)}} R_J^{(n)} \rightarrow R_I^{(n)}$$

which is precisely $C_I^{(n)}$, proving (2) and (3). $\square$

Corollary 4.2. *The category $\text{Mod}(Z_n(A))$ is equivalent to the category $\text{Rep}(Q_{A, n-N+1})$ where $Q_{A, n-N+1}$ is the quiver with relations which has a vertex x_I for every admissible $I \subset \{1, \dots, n-N+1\}$ and arrows $x_{IJ} : x_I \rightarrow x_J$ for every $I \subset J$. The relations are given by $x_{JK}x_{IJ} = x_{IK}$. If $n-N+1 \leq 0$ then $Q_{A, n-N+1}$ is a single vertex $x_\emptyset$ and no arrows.*

Proof. This is an easy consequence of the description of the indecomposable projectives $Z_n(A)$ -representations in Proposition 4.1. The functor realizing the stated equivalence is $\text{Hom}_{Z_n(A)}(\bigoplus_{I \text{ adm.}} R_I, -)$. If P_I is the projective $Q_{A, n-N+1}$ representation corresponding to the vertex I then the inverse equivalence sends P_I to R_I . $\square$

For use below we denote the inverse equivalence alluded to in the above proof by $\mathcal{F}_A^\circ$.

If we apply Theorem 2.1.1 with $\mathcal{C} = \text{CoMod}(\underline{\text{end}}(A)_n)$ then we obtain an exact functor

$$\mathcal{F}_A : \text{Cube}_{n-N+1} \rightarrow \text{CoMod}(\underline{\text{end}}(A)_n)$$

which sends $P_I^{(n-N+1)}$ to $R_I^{(n-N+1)}$ and which extends to an exact monoidal functor

$$\mathcal{F}_A : \text{Cube}_\bullet \rightarrow \text{CoMod}(\underline{\text{end}}(A))$$

The functor $\mathcal{F}_A$ is an enhancement of the functor F_A we used before as it fits in the following commutative diagram

$$\begin{array}{ccc} & \text{CoMod}(\underline{\text{end}}(A)_n) & \\ \mathcal{F}_A \nearrow & & \searrow \text{forget} \\ \text{Cube}_{n-N+1} & \xrightarrow{F_A} & \text{Vect} \end{array}$$

Proposition 4.3. *The functor $\mathcal{F}_A$ defines an equivalence*

$$\bar{\mathcal{F}}_A : \text{Cube}_\bullet / \mathcal{S}_A \cong \text{CoMod}(\underline{\text{end}}(A))$$

where $\mathcal{S}_A$ is the localizing subcategory of $\text{Cube}_\bullet$ generated by the simples $S_I^{(l)}$ with I not admissible.

Proof. For conciseness we drop some superscripts in the notations below, when they are clear from the context. We consider the following diagram

$$(4.2) \quad \begin{array}{ccc} & \text{Rep}(Q_{A,n-N+1}) & \\ \text{Res} \nearrow & & \searrow \mathcal{F}_A^\circ \\ \text{Cube}_{n-N+1} & \xrightarrow{\mathcal{F}_A} & \text{Mod}(Z_n(A)) \end{array}$$

Ind

Here Res is the restriction functor $\text{Cube}_n = \text{Rep}(Q_n) \rightarrow \text{Rep}(Q_{A,n})$ and Ind is the left adjoint to Res .

We first claim $\mathcal{F}_A^\circ = \mathcal{F}_A \circ \text{Ind}$. To see this note that all functors are right exact and commute with direct sums, so it is sufficient to show that they take the same value on projectives. If I is admissible then we find $(\mathcal{F}_A \circ \text{Ind})(P_I) = \mathcal{F}_A(P_I) = R_I$ and we find the same value for $\mathcal{F}_A^\circ(P_I)$.

Now we claim $\mathcal{F}_A = \mathcal{F}_A^\circ \circ \text{Res}$. Let $M \in \text{Cube}_{n-N+1}$. It follows that

$$(\mathcal{F}_A^\circ \circ \text{Res})(M) = \mathcal{F}_A(\text{Ind} \circ \text{Res}(M))$$

The canonical map

$$\text{Ind} \circ \text{Res}(M) \rightarrow M$$

is surjective and has its kernel in $\mathcal{S}_A$. Hence since $\mathcal{F}_A$ is exact

$$\mathcal{F}_A(\text{Ind} \circ \text{Res}(M)) = \mathcal{F}_A(M)$$

which implies our claim. Thus diagram (4.2) is commutative in the two possible senses.

Put $\mathcal{S}_{A,n} = \mathcal{S}_A \cap \text{Cube}_n$. Then $\mathcal{S}_{A,n}$ lies in the kernel of Res and hence in the kernel of $\mathcal{F}_A$. Furthermore Res induces an equivalence $\text{Cube}_n / \mathcal{S}_{A,n} \cong \text{Rep}(Q_{A,n})$. Hence we obtain a commutative diagram

$$(4.3) \quad \begin{array}{ccc} & \text{Rep}(Q_{A,n-N+1}) & \\ \text{Res} \nearrow & & \searrow \mathcal{F}_A^\circ \\ \text{Cube}_{n-N+1} / \mathcal{S}_{A,n-N+1} & \xrightarrow{\bar{\mathcal{F}}_A} & \text{Mod}(Z_n(A)) \end{array}$$

$\cong$

Since $\mathcal{F}_A^\circ$ is an equivalence, the same holds for $\bar{\mathcal{F}}_A$. □

Corollary 4.4. *Assume that A satisfies the extra condition (besides being distributive). Then the monoidal category $\text{CoMod}(\underline{\text{end}}(A))$ is entirely determined by the numbers $(\dim J_n)_n$. If $N = 2$ then $\text{CoMod}(\underline{\text{end}}(A))$ is entirely determined by the Hilbert series of A .*

Proof. It follows from Proposition 4.3 that to describe the category $\text{CoMod}(\underline{\text{end}}(A))$ it suffices to know which $S_I^{(l)}$ are mapped to zero under $\mathcal{F}_A$ or equivalently under F_A .

To this end it is sufficient to compute the dimension of $F_A S_I^{(l)}$. First we observe that $S_I^{(l)}$ has a finite projective resolution

$$\cdots \rightarrow \bigoplus_{\substack{J \supset I \\ |J-I|=2}} P_J^{(l)} \rightarrow \bigoplus_{\substack{J \supset I \\ |J-I|=1}} P_J^{(l)} \rightarrow S_I^{(l)} \rightarrow 0$$

Hence to know $\dim F_A S_I^{(l)}$ it is sufficient to know $\dim F_A P_I^{(l)}$.

Since A satisfies the extra condition we have $\dim F_A P_I^{(l)} = \dim F_A P_{\bar{I}}^{(l)}$ where $\bar{I}$ is obtained from I by filling up all holes of size $\leq N-2$. Thus we may assume that I has no holes of size $\leq N-2$.

If on the other hand I contains a hole of size $\geq N-1$ or else does not contain 1 or l then we may write $P_I^{(l)}$ as a tensor product $P_{I_1}^{(l_1)} \otimes P_{I_2}^{(l_2)}$ for suitable l_1, l_2, I_1, I_2 . Hence we may reduce to the case where $I = \{1, \dots, l\}$. In that case $F_A P_I^{(l)} = J_l$. This finishes the proof for general N .

If $N = 2$ then it follows from the Koszul complex that the sets of numbers $(\dim J_n)_n$ and $(\dim A_n)_n$ determine each other. $\square$

5. THE SYMMETRIC ALGEBRA

In this section we assume that the ground field k is algebraically closed of characteristic zero. Below A will be the symmetric algebra $SV = TV/(\wedge^2 V)$ of a vector space V of dimension n . As SV is a quadratic algebra we will have $N = 2$ in this section.

To simplify things conceptually we will equip V with an explicit basis $(x_i)_i$ as in §3 and we put $z_i^j = x_j^* \otimes x_i$. Let $\mathcal{O} = k[(z_i^j)_{i,j}] = S(\text{End}(V)^*)$ be the coordinate ring of $n \times n$ -matrices. Then $\mathcal{O}$ is a bialgebra which coacts on SV with the same formulas as (3.1)(3.2). Furthermore sending $z_i^j \mapsto z_i^j$ defines a bialgebra homomorphism $B : \underline{\text{end}}(A) \rightarrow \mathcal{O}$ compatible with the coactions on SV . Hence we obtain a corresponding monoidal exact functor

$$B_* : \text{CoMod}(\underline{\text{end}}(A)) \rightarrow \text{CoMod}(\mathcal{O})$$

which on the underlying vector spaces is the identity.

The bialgebra $\mathcal{O}$ is semi-simple and its irreducible representations are indexed by partitions of at most n rows. For more details we refer to the excellent text book [11]. If λ is a partition of m with conjugate partition λ' then the associated irreducible $\mathcal{O}$ (co)representation is

$$S^\lambda V = \text{im} \left(\bigotimes_i \wedge^{\lambda'_i} V \xrightarrow{\alpha} V^{\otimes n} \xrightarrow{\beta} \bigotimes_j S^{\lambda_j} V \right)$$

The precise form of the (anti-)symmetrization maps α and β is derived from a labeling by the numbers $1, \dots, m$ of the Young diagram associated to λ (which

does not have to yield a standard tableau). The representation $S^\lambda V$ is independent of this labeling.

For example if $\lambda = [221]$ then a corresponding labeled Young diagram is

1	2
3	4
5	

The corresponding irreducible $\mathcal{O}$ -representation $S^\lambda V$ is given by

$$\text{im}(\wedge^3 V \otimes \wedge^2 V \xrightarrow{\alpha} V^{\otimes 5} \xrightarrow{\beta} S^2 V \otimes S^2 V \otimes V)$$

The image of α consists of the tensors which are anti-symmetric in the places (1, 3, 5) and (2, 4). Likewise the map β symmetrizes a tensor in the places (1, 2), (3, 4) (and 5 but this has no effect of course).

The construction of $S^\lambda V$ works also when λ is replaced by a skew diagram λ/μ , i.e. a difference of two partitions λ, μ . In that case we have

$$S^{\lambda/\mu} V = \text{im} \left(\bigotimes_i \wedge^{\lambda'_i - \mu'_i} V \xrightarrow{\alpha} V^{\otimes n} \xrightarrow{\beta} \bigotimes_j S^{\lambda_j - \mu_j} V \right)$$

For example if λ/μ is the skew (labeled) diagram

	1
2	3
4	

then

$$S^{\lambda/\mu} V = \text{im}(\wedge^2 V \otimes \wedge^2 V \rightarrow V^{\otimes 4} \rightarrow V \otimes S^2 V \otimes V)$$

In contrast to $S^\lambda V$ the representation $S^{\lambda/\mu} V$ is in general not irreducible but its decomposition can be described in terms of the well-known Littlewood-Richardson coefficients.

$$S^{\lambda/\mu} V = \sum_{\nu} (S^\nu)^{\oplus N_{\lambda\mu\nu}}$$


The Littlewood-Richardson coefficients can be computed by means of a combinatorial recipe which consists in counting certain admissible labelings of the boxes of λ/μ .

The following is the main observation of this section.

Proposition 5.1. *Assume*

$$I = \{1, \dots, a, \widehat{a+1}, \dots, \widehat{a+b}, a+b+1, \dots, a+b+c, \dots\} \subset \{1, \dots, m\}$$

Then $B_*(\mathcal{F}_A(S_I^{(m)})) = S^{\lambda/\mu} V$ where λ/μ is the rim hook with $m+1$ boxes:



(5.1)

such that the column labeled by “1” has $a + 1$ boxes, the row labeled by “2” has $b + 1$ -boxes, the row labeled by “3” has $c + 1$ -boxes, etc. . . .

This result remains valid if I has instead the form

$$I = \{\hat{1}, \dots, \hat{b}, b + 1, \dots, b + c, \dots\}$$

In that case we simply think of a as being zero. This means that the column labeled 1 contains only one box and thus is contained in the row labeled 2.

Proof. It is easy to see that

$$\begin{aligned} B_*(\mathcal{F}_A(S_{\{1, \dots, l\}}^{(l)})) &= B_*(\wedge^l V) = \wedge^l V \\ B_*(\mathcal{F}_A(S_{\emptyset}^{(l)})) &= B_*(S^l V) = S^l V \end{aligned}$$

An arbitrary $S_I^{(m)}$ has a presentation

$$\bigoplus_{i \notin I} P_{I \cup \{i\}}^{(m)} \rightarrow P_I^{(m)} \rightarrow S_I^{(m)} \rightarrow 0$$

Applying $\mathcal{F}_A$ and using distributivity we get

$$\mathcal{F}_A S_I^{(m)} = R_I^{(m+1)} \Big/ \left(R_I^{(m+1)} \cap \sum_{i \notin I} R_i^{(m+1)} \right)$$

Alternatively

$$\mathcal{F}_A S_I^{(m)} = \text{im} \left(R_I^{(m+1)} \rightarrow V^{\otimes m+1} \rightarrow V^{\otimes m+1} \Big/ \sum_{i \notin I} R_i^{(m+1)} \right)$$

We have

$$\begin{aligned} R_I^{(m+1)} &= \wedge^{a+1} V \otimes \wedge^{c+1} \otimes \dots \\ V^{\otimes m+1} \Big/ \sum_{i \notin I} R_i^{(m+1)} &= S^{b+1} V \otimes S^{d+1} V \otimes \dots \end{aligned}$$

Thus $\mathcal{F}_A S_I^{(m)}$ is $S^{\lambda/\mu} V$ where λ/μ is the diagram (5.1) numbered with the numbers $1, \dots, m + 1$ starting from the top right and ending at the bottom left (thus this is not a standard tableau). $\square$

6. SOME EXAMPLES OF DISTRIBUTIVE ALGEBRAS

In this section we describe some algebras which we could show to be distributive.

6.1. Algebras derived from Koszul algebras. We have the following easy general result.

Proposition 6.1.1. *Let $A = TV/(R)$ be a quadratic Koszul algebra and let $R' \in \mathcal{L}_N(A)$. Then $A' = TV/(R')$ is a distributive algebra. If specifically we take*

$$(6.1) \quad R' = \bigcap_{i=1}^{N-1} V^{\otimes i-1} \otimes R \otimes V^{N-i-1}$$

then A' also satisfies the extra condition and hence is N -Koszul.

Proof. The distributivity of A' follows immediately from the distributivity of A . To check the extra condition we have to verify for $1 \leq l \leq p$ with $p \leq N - 1$

$$(6.2) \quad \left(\bigcap_{i=1}^{N-1} V^{\otimes i-1} \otimes R \otimes V^{N+(p-1)-i-1} \right) \cap \left(\bigcap_{i=1}^{N-1} V^{\otimes (p-1)+i-1} \otimes R \otimes V^{N-i-1} \right) \\ \subset \bigcap_{i=1}^{N-1} V^{\otimes (l-1)+i-1} \otimes R \otimes V^{N-(l-1)-i-1}$$

The lefthand side of this equation is equal to

$$\bigcap_{j=1}^{p+N-2} V^{\otimes j-1} \otimes R \otimes V^{N+p-2-j}$$

We claim that each of the terms on the righthand side of (6.2) appears in this intersection. To prove this we have to find for any $i = 1, \dots, N-1$ a $j \in \{1, \dots, p+N-2\}$ such that $j-1 = l+i-2$. Thus we have to take $j = l+i-1$. One verifies that indeed $1 \leq l+i-1 \leq p+N-2$. $\square$

Example 6.1.2. Take $R = \wedge^2 V$ and R' as (6.1). Then $R' = \wedge^N V$ and hence $A' = TV/(\wedge^N V)$ is the N -generalization of the symmetric algebra introduced in [6]. By Proposition 6.1.1 we see that A' is distributive and satisfies the extra condition (and hence is N -Koszul by (2.4)). These facts were proved by Berger in [6] using “confluence”. See below.

6.2. Confluence. For $N > 2$ the only tool to establish distributivity of $TV/(R)$ with $R \subset V^{\otimes N}$ seems to be “confluence” [6, 8]. For the benefit of the reader we give a quick introduction to this concept.

Let W be a finite dimensional vector space equipped with a fixed totally ordered basis X . If $S \subset W$ is a subvector space then an element x of X is called non-reduced with respect to S if S contains an element of the form $x - \sum_{y < x} c_y y$. We denote the set of such non-reduced monomials by $\text{NRed}(S)$. The following is easy to see.

Lemma 6.2.1. (1) $\dim S = |\text{NRed}(S)|$.
(2) If $S \subset R$ then $\text{NRed}(S) \subset \text{NRed}(R)$.

Thus $\text{NRed}(-)$ is an order preserving map from the lattice of subvector spaces of W to the lattice of subsets of X .

From the fact that $\text{NRed}(-)$ is order preserving we immediately deduce

$$(6.3) \quad \text{NRed}(R \cap S) \subset \text{NRed}(R) \cap \text{NRed}(S)$$

$$(6.4) \quad \text{NRed}(R + S) \supset \text{NRed}(R) \cup \text{NRed}(S)$$

The following result was proved by Berger [5].

Lemma 6.2.2. If one of these inclusions is an equality then so is the other.

Definition 6.2.3. We say that $R, S \subset W$ are confluent if one (and hence both) of the inclusions (6.3)(6.4) is an equality.

The following is one of the main fact about confluence [5].

Theorem 6.2.4. If $\mathcal{R} = \{(R_i)_i\}$ is a collection of pairwise confluent subspaces of W then $\text{NRed}(-)$ defines a isomorphism between the lattice $\mathcal{L}$ of subspaces generated by R_i and the lattice of subsets of X generated by $\text{NRed}(R_i)$. In particular $\mathcal{L}$ is distributive.

A typical application of this concept is the following. Let $A = TV/(R)$ with $R \subset V^{\otimes N}$. Let X be a totally ordered basis for V . We equip $V^{\otimes n}$ with the basis $X^{\times n}$, ordered lexicographically. Then we say that A is confluent (with respect to X) if for all n and all i the subspaces $R_i^{(n)}$ of $V^{\otimes n}$ are pairwise confluent. Using Theorem 6.2.4 we get

$$\text{confluent} \Rightarrow \text{distributive}$$

The following was observed by Berger [5].

Proposition 6.2.5. *In order for A to be confluent with respect to X it is necessary and sufficient that $R \otimes V^{\otimes i}$ and $V^{\otimes i} \otimes R$ are confluent inside $V^{\otimes(i+N)}$ for $i = 1, \dots, N-1$.*

6.3. The (deformed) Yang-Mills algebras. In this section we assume that k has characteristic zero. The Yang-Mills algebras and their deformations were introduced by Connes and Dubois-Violette in [9, 10]. After a short reminder of how they are constructed we will show that they are distributive.

The i 'th cyclic derivative on $F = k\langle x_1, \dots, x_n \rangle$ is a k -linear map

$$\frac{\circ \partial}{\partial x_i} : F/[F, F] \rightarrow F$$

which on monomials is defined by

$$\frac{\circ \partial m}{\partial x_i} = \sum_{m=ux_i v} vu$$

Assume that V is an n -dimensional vector space equipped with a basis $(x_i)_i$. To avoid trivialities we assume $n \geq 2$. If $w \in TV/[TV, TV]$ is homogeneous then the Jacobian algebra associated to w is the algebra $A = TV/(R)$ where R is the vectorspace spanned by the cyclic derivatives of w . It is easy to see that R and hence A does not depend on the choice of the basis $(x_i)_i$.

One has

$$\bar{w} \stackrel{\text{def}}{=} \sum_i x_i \frac{\circ \partial w}{\partial x_i} = \sum_i \frac{\circ \partial w}{\partial x_i} x_i \in R \otimes V \cap V \otimes R$$

The map $w \mapsto \bar{w}$ gives an isomorphism between $(F/[F, F])_n$ and the cyclically invariant elements of $F_n = V^{\otimes n}$.

Assume now that V is equipped with a non-degenerate symmetric bilinear form $(-, -)$ and put $g_{ij} = (x_i, x_j)$. The inverse of the matrix $(g_{ij})_{ij}$ is denoted by $(g^{ij})_{ij}$. Let $O(V)$ be the corresponding orthogonal group. Then one has the following result

Lemma 6.3.1. *The space $(TV/[TV, TV])_4^{O(V)}$ is two-dimensional and is spanned by the elements*

$$w_1 = \sum_{i,j,p,q} g^{ip} g^{jq} [x_i, x_j] [x_p, x_q]$$

$$w_2 = \left(\sum_{i,j} g^{ij} x_i x_j \right)^2$$

Proof. To prove this we may assume that $k = \bar{k}$ and that $(x_i)_i$ is an orthogonal basis for V . In other words $g_{ij} = \delta_{ij}$. We first consider $(V^{\otimes 4})^{O(V)}$. The first

fundamental theorem of invariant theory for the orthogonal groups implies that $(V^{\otimes 4})^{O(V)}$ is spanned by

$$\begin{aligned} s_1 &= \sum_{i,j} x_i x_i x_j x_j \\ s_2 &= \sum_{i,j} x_i x_j x_i x_j \\ s_3 &= \sum_{i,j} x_i x_j x_j x_i \end{aligned}$$

Since $n \geq 2$ it is clear that s_1, s_2, s_3 are linearly independent. Hence they form a basis for $(V^{\otimes 4})^{O(V)}$.

Since $O(V)$ is reductive the quotient map $(V^{\otimes 4})^{O(V)} \rightarrow (TV/[TV, TV])_4^{O(V)}$ is surjective and hence $(TV/[TV, TV])_4^{O(V)}$ is spanned by s_1, s_2, s_3 . It is clear that s_1 and s_3 are equal modulo commutators but s_1 and s_2 remain independent. We now use the fact that $w_2 = s_1$ and $w_1 = 2(s_2 - s_3)$. $\square$

By definition the deformed Yang-Mills algebra A_λ ($\lambda \in k$) is the Jacobian algebra for the potential $w_1 + \lambda w_2$. The equations of A_λ have the form

$$\sum_{j,p} g^{jp} ([x_j, [x_i, x_p]] + \lambda \{x_i, x_j x_p\})$$

where $\{a, b\} = ab + ba$. The ordinary Yang-Mills algebra is A_0 .

Theorem 6.3.2. *The deformed Yang-Mills algebra A_λ is distributive and satisfies the extra condition.*

Proof. We may clearly assume that the ground field is algebraically closed. Then we may assume that g^{ij} has the following form

$$g^{ij} = \begin{cases} 1 & \text{if } i + j = n \\ 0 & \text{otherwise} \end{cases}$$

Put $\bar{i} = n - i$. The equations of A_λ now have the form

$$\sum_j [x_j, [x_i, x_j]] + \lambda \{x_i, x_j x_j\}$$

Thus we get

$$\text{NRed}(R) = \{x_n x_i x_1 \mid 1 \leq i \leq n\}$$

We claim that A_λ is confluent. By Proposition 6.2.5 we must check the equalities

$$\text{NRed}(R_1^{(n)} \cap R_{n-2}^{(n)}) = \text{NRed}(R_1^{(n)}) \cap \text{NRed}(R_{n-2}^{(n)})$$

for $n = 4, 5$. We already know that the sought equalities are inclusions of the type “ $\subset$ ”. Hence it is sufficient to prove

$$(6.5) \quad \dim R_1^{(n)} \cap R_{n-2}^{(n)} \geq |\text{NRed}(R_1^{(n)}) \cap \text{NRed}(R_{n-2}^{(n)})|$$

Since $\text{NRed}(R_1^{(5)}) \cap \text{NRed}(R_3^{(5)}) = \emptyset$ there is nothing to prove for $n = 5$.

On the other hand $\text{NRed}(R_1^{(4)}) \cap \text{NRed}(R_2^{(4)}) = \{x_n x_n x_1 x_1\}$ and $\bar{w}_\lambda \in R_1^{(4)} \cap R_2^{(4)}$. Thus $\dim R_1^{(4)} \cap R_2^{(4)} \geq 1$ and so (6.5) holds for $n = 4$ as well.

To check the extra condition, using Theorem 6.2.4 we must verify

$$\text{NRed}(R_1^{(5)}) \cap \text{NRed}(R_3^{(5)}) \subset \text{NRed}(R_2^{(5)})$$

As the lefthand side is empty there is nothing to prove. $\square$

Using (2.4) we recover the following result proved in [9, 10].

Corollary 6.3.3. *The deformed Yang-Mills algebra A_λ is 3-Koszul.*

6.4. Three dimensional cubic Artin-Schelter regular algebras. Recall [1] that an AS-regular algebra is a graded k -algebra $A = k + A_1 + A_2 + \dots$ satisfying the following conditions

- (1) $\dim A_i$ is bounded by a polynomial.
- (2) The projective dimension of k is finite.
- (3) There is exactly one i for which $\text{Ext}_A^i(k, A)$ is non-vanishing and for this i we have $\dim \text{Ext}_A^i(k, A) = 1$.

Three dimensional regular algebras generated in degree one were classified in [1, 2, 3] and in general in [16, 17]. It was discovered that they are intimately connected to plane elliptic curves.

There are two possibilities for a three dimensional regular algebra A generated in degree one.

- (1) A is defined by three generators satisfying three quadratic relations (the “quadratic case”).
- (2) A is defined by two generators satisfying two cubic relations (“the cubic case”).

In [1] it is shown that all 3-dimensional regular algebras are obtained by specialization from a number “generic” regular algebras. These generic regular algebras depend on at most two parameters.

Quadratic three dimensional AS-regular algebras are Koszul and hence distributive. Cubic three dimensional AS-regular algebras are 3-Koszul so one would expect it should be easy to verify distributivity for them. Nonetheless we have not found a clean way to establish this.

Below we discuss so-called Type A algebras by using a generalization of the notion of I -type introduced in [18]. It is likely that the other types can be handled in a similar way but we have not carried out the required verifications.

Type A algebras are associated to a triple $(E, \sigma, \mathcal{L})$ where E is a smooth elliptic curve, σ is a translation and $\mathcal{L}$ is a line bundle of degree two on E . Put $V = H^0(E, \mathcal{L})$ and

$$R = \ker(H^0(E, \mathcal{L})^{\otimes 3} \xrightarrow{a \otimes b \otimes c \mapsto ab^\sigma c^{\sigma^2}} H^0(E, \mathcal{L} \otimes \mathcal{L}^\sigma \otimes \mathcal{L}^{\sigma^2}))$$

(where $(-)^{\sigma} = \sigma^*$). Then the cubic three-dimensional AS-regular algebra associated to $(E, \sigma, \mathcal{L})$ is given by $TV/(R)$.

We prove

Proposition 6.4.1. *Let A be a cubic Type A regular algebra. Then A is distributive.*

It follows easily from [1, Thm 6.11] that a cubic Type A regular algebra is not confluent in the sense §6.2 for any choice of basis X . Our arguments still use confluence but in a more sophisticated way.

We need some preparatory work. There is a well-defined map

$$\phi : A_n \rightarrow H^0(E, \mathcal{L} \otimes \cdots \otimes \mathcal{L}^{\sigma^{n-1}}) : \overline{a_0 \otimes \cdots \otimes a_n} \mapsto a_0 a_1^\sigma \cdots a_{n-1}^{\sigma^{n-1}}$$

If $f \in A_n$ then we write (f) for the divisor of the image of $\phi(f)$ in $H^0(E, \mathcal{L} \otimes \cdots \otimes \mathcal{L}^{\sigma^{n-1}})$. If $f \in A_m, g \in A_n$ then

$$(fg) = (\phi(fg)) = (\phi(f)\phi(g)^{\sigma^m}) = (f) + \sigma^{-m}(g)$$

Let x, y be a basis of $V = H^0(E, \mathcal{L})$. Put $(x) = P + P', (y) = Q + Q'$. We write x_n, y_n for arbitrary non-zero elements in V with divisors $\sigma^n P + \sigma^{-n} P', \sigma^n Q + \sigma^{-n} Q'$.

We will say that P, P', Q, Q' are generic if they have pairwise disjoint orbits.

Lemma 6.4.2. *Assume that P, P', Q, Q' are generic and $\sigma^4 \neq \text{id}$. Then there are non-zero scalars $\alpha, \beta, \gamma, \delta$ such that R has a basis given by*

$$(6.6) \quad \begin{aligned} & y_0 x_1 x_{-2} - \alpha x_0 x_{-3} y_2 - \beta x_0 y_1 x_{-2} \\ & y_0 y_{-3} x_2 - \gamma x_0 y_1 y_{-2} - \delta y_0 x_1 y_{-2} \end{aligned}$$

Proof. We need to prove that the elements (6.6) are linearly independent and contained in R . We first show that they are linearly independent. Assume this is not the case. I.e. there are scalars λ, μ , not both zero such that

$$\lambda(y_0 x_1 x_{-2} - \alpha x_0 x_{-3} y_2 - \beta x_0 y_1 x_{-2}) + \mu(y_0 y_{-3} x_2 - \gamma x_0 y_1 y_{-2} - \delta y_0 x_1 y_{-2}) = 0$$

in $V^{\otimes 3}$. Collecting the terms starting with x_0

$$-\lambda \alpha x_{-3} y_2 - \lambda \beta y_1 x_{-2} - \mu \gamma y_1 y_{-2} = 0$$

By our genericity assumption x_{-3} and y_1 are linearly independent. Thus we get

$$\begin{aligned} -\lambda \alpha y_2 &= 0 \\ -\lambda \beta x_2 - \mu \gamma y_{-2} &= 0 \end{aligned}$$

From the first equation we deduce $\lambda = 0$ and from the second $\mu = 0$.

Now we show that the elements (6.6) are indeed contained in R (for suitable choices of scalars). To this end we have to show that the element

$$y_0 x_1^\sigma x_{-2}^{\sigma^2}$$

of $H^0(E, \mathcal{L} \otimes \mathcal{L}^\sigma \otimes \mathcal{L}^{\sigma^2})$ is a linear combination with non-zero coefficients of

$$x_0 x_{-3} y_2^{\sigma^2}, \quad x_0 y_1^\sigma x_{-2}^{\sigma^2}$$

and similarly the element

$$y_0 y_{-3}^\sigma x_2^{\sigma^2}$$

is a linear combination with non-zero coefficients of

$$x_0 y_1^\sigma y_{-2}^{\sigma^2}, \quad y_0 x_1^\sigma y_{-2}^{\sigma^2}$$

We compute the divisors of all these elements

$$\begin{aligned}
y_0 x_1^\sigma x_{-2}^{\sigma^2} : & \quad Q + Q' + P + \sigma^{-2} P' + \sigma^{-4} P + P' \\
x_0 x_{-3}^\sigma y_2^{\sigma^2} : & \quad P + P' + \sigma^{-4} P + \sigma^2 P' + Q + \sigma^{-4} Q' \\
x_0 y_1^\sigma x_{-2}^{\sigma^2} : & \quad P + P' + Q + \sigma^{-2} Q' + \sigma^{-4} P + P' \\
y_0 y_{-3}^\sigma x_2^{\sigma^2} : & \quad Q + Q' + \sigma^{-4} Q + \sigma^2 Q' + P + \sigma^{-4} P' \\
x_0 y_1^\sigma y_{-2}^{\sigma^2} : & \quad P + P' + Q + \sigma^{-2} Q' + \sigma^{-4} Q + Q' \\
y_0 x_1^\sigma y_{-2}^{\sigma^2} : & \quad Q + Q' + P + \sigma^{-2} P' + \sigma^{-4} Q + Q'
\end{aligned}$$

All these divisors are different so these elements are pairwise linearly independent. We also find

$$y_0 x_1^\sigma x_{-2}^{\sigma^2}, x_0 x_{-3}^\sigma y_2^{\sigma^2}, x_0 y_1^\sigma x_{-2}^{\sigma^2} \in H^0(E, \mathcal{L} \otimes \mathcal{L}^\sigma \otimes \mathcal{L}^{\sigma^2}(-P - P' - Q - \sigma^{-4} P))$$

By Riemann Roch the righthand side is a two dimensional vector space. Hence $y_0 x_1^\sigma x_{-2}^{\sigma^2}$, $x_0 x_{-3}^\sigma y_2^{\sigma^2}$, $x_0 y_1^\sigma x_{-2}^{\sigma^2}$ are linearly dependent. Since they are not scalar multiples of each other the first one is indeed a linear combination of the other two and the coefficients are non-zero.

Similarly we find


$$y_0 y_{-3}^\sigma x_2^{\sigma^2}, x_0 y_1^\sigma y_{-2}^{\sigma^2}, y_0 x_1^\sigma y_{-2}^{\sigma^2} \in H^0(E, \mathcal{L} \otimes \mathcal{L}^\sigma \otimes \mathcal{L}^{\sigma^2}(-P - Q - Q' - \sigma^{-4} Q))$$

With the same reasoning as above we deduce that $y_0 y_{-3}^\sigma x_2^{\sigma^2}$ is a linear combination of $x_0 y_1^\sigma y_{-2}^{\sigma^2}$, $y_0 x_1^\sigma y_{-2}^{\sigma^2}$ with non-zero coefficients. $\square$


Proof of Proposition 6.4.1. We will give the proof in the case that $\sigma^4 \neq \text{id}$. If $\sigma^4 = \text{id}$ then A is a so-called “linear” algebra which can be easily analyzed directly (see [2, Prop. 7.4]).

We may clearly assume that our base field is so large that we can select generic P, P', Q, Q' as in the discussion above.

Let us grade the monomials in x, y by putting $\deg x = (1, 0)$, $\deg y = (0, 1)$. For monomials μ in x, y and $(a, b) \in \mathbb{Z}^2$ we define $v_{a,b}(\mu) \in TV$ by constructing a path in the grid below starting at (a, b) and ending at $(a, b) + |\mu|$ where the variables x, y indicate which branch we should take at each vertex.



One square looks like



Thus for example

$$v_{0,0}(xyyx) = x_0y_1y_{-2}x_{-1}$$

It is clear that $v_{a,b}(\mu)_\mu$ is a basis for TV for any a, b . We extend $v_{a,b}$ linearly to an isomorphism of graded vector spaces $TV \rightarrow TV$. If $(a, b) = (0, 0)$ then we write $v(\mu) = v_{a,b}(\mu)$.

We claim that $V^{\otimes a} \otimes R \otimes V^{\otimes b}$ has a basis of the form

$$(6.8) \quad v(\mu y x x \mu') - \alpha_\mu v(\mu x x y \mu') - \beta_\mu v(\mu x y x \mu')$$

$$(6.9) \quad v(\mu y y x \mu') - \gamma_\mu v(\mu x y y \mu') - \delta_\mu v(\mu y x y \mu')$$

where μ, μ' run through the monomials of length a, b and $\alpha_\mu, \beta_\mu, \gamma_\mu, \delta_\mu$ are scalars depending on μ .

By Lemma 6.4.2 we find that R has a basis of the type

$$r'_{m,n} = y_n x_{m+1} x_{m-2} - \alpha_{m,n} x_m x_{m-3} y_{n+2} - \beta_{m,n} x_m y_{n+1} x_{m-2}$$

$$s'_{m,n} = y_n y_{n-3} x_{m+2} - \gamma_{m,n} x_m y_{n+1} y_{n-2} - \delta_{m,n} y_n x_{m+1} y_{n-2}$$

for each m, n . Put

$$r_{m,n} = y x x - \alpha_{m,n} x x y - \beta_{m,n} x y x$$

$$s_{m,n} = y y x - \gamma_{m,n} x y y - \delta_{m,n} y x y$$

If $(m, n) = (-3p + q, -3q + p)$ then

$$v_{p,q}(r_{m,n}) = r'_{m,n}$$

$$v_{p,q}(s_{m,n}) = s'_{m,n}$$

Assume now that μ, μ' are monomials of degree $(p, q), (p', q')$ respectively with $p + q = a, p' + q' = b$. Put $(m, n) = (-3p + q, -3q + p)$. Then

$$v(\mu r_{m,n} \mu') = v(\mu) v_{p,q}(r_{m,n}) v_{p+2,q+1}(\mu')$$

$$v(\mu s_{m,n} \mu') = v(\mu) v_{p,q}(s_{m,n}) v_{p+1,q+2}(\mu')$$

Thus $v(\mu r_{m,n} \mu'), v(\mu s_{m,n} \mu') \in V^{\otimes a} \otimes R \otimes V^{\otimes b}$. It is clear that the elements $\mu r_{m,n} \mu', \mu s_{m,n} \mu'$ for varying μ, μ' are all linearly independent and their number is equal to the dimension of $V^{\otimes a} \otimes R \otimes V^{\otimes b}$. So they yield a basis for $V^{\otimes a} \otimes R \otimes V^{\otimes b}$ after applying $v(-)$. This finishes the proof of our claim.

We now claim that the $(R_a^{(n)})_a$ are pairwise confluent inside $V^{\otimes n}$ when the latter is equipped with the ordered basis $v(\mu)_\mu$ where μ runs through the lexicographically ordered set of monomials of length n . The proposition then follows from Theorem 6.2.4.

We compute for $1 \leq a < b \leq n - 3$

$$|\mathrm{NRed}(R_a^{(n)}) \cap \mathrm{NRed}(R_b^{(n)})| = \begin{cases} 2^{n-4} & \text{if } b = a + 1 \\ 0 & \text{if } b = a + 2 \\ 2^{n-4} & \text{if } b \geq a + 3 \end{cases}$$

If $b \geq a + 3$ then

$$\dim R_a^{(n)} \cap R_b^{(n)} = 2^{n-4}$$

Hence the only statement that needs to be shown is

$$\dim R_a^{(n)} \cap R_{a+1}^{(n)} = 2^{n-4}$$

This follows from the fact that for a cubic three-dimensional AS-regular algebra we have $\dim(R \otimes V \cap V \otimes R) = 1$ [1, Prop. (2.4)].

REFERENCES

- [1] M. Artin and W. Schelter, *Graded algebras of global dimension 3*, Adv. in Math. **66** (1987), 171–216.
- [2] M. Artin, J. Tate, and M. Van den Bergh, *Some algebras associated to automorphisms of elliptic curves*, The Grothendieck Festschrift, vol. 1, Birkhäuser, 1990, pp. 33–85.
- [3] ———, *Modules over regular algebras of dimension 3*, Invent. Math. **106** (1991), 335–388.
- [4] J. Backelin and R. Fröberg, *Koszul algebras, Veronese subrings and rings with linear resolutions*, Rev. Roumaine Math. Pures Appl. **30** (1985), no. 2, 85–97.
- [5] R. Berger, *Confluence and Koszulity*, J. Algebra **201** (1998), no. 1, 243–283.
- [6] ———, *Koszulity for nonquadratic algebras*, J. Algebra **239** (2001), no. 2, 705–734.
- [7] R. Berger, M. Dubois-Violette, and M. Wambst, *Homogeneous algebras*, J. Algebra **261** (2003), no. 1, 172–185.
- [8] G. M. Bergman, *The diamond lemma for ring theory*, Adv. in Math. **29** (1978), no. 2, 178–218.
- [9] A. Connes and M. Dubois-Violette, *Yang-Mills algebra*, Lett. Math. Phys. **61** (2002), no. 2, 149–158.
- [10] ———, *Yang-Mills and some related algebras*, Rigorous quantum field theory (Basel), Progr. Math., vol. 251, Birkhäuser, Basel, 2007, pp. 65–78.
- [11] W. Fulton and J. Harris, *Representation theory*, Graduate Texts in Mathematics, vol. 129, Springer-Verlag, New York, 1991, A first course, Readings in Mathematics.
- [12] P. H. Hai, B. Kriegk, and M. Lorenz, *N-homogeneous superalgebras*, J. Noncommut. Geom. **2** (2008), no. 1, 1–51.
- [13] P. H. Hai and M. Lorenz, *Koszul algebras and the quantum MacMahon master theorem*, Bull. Lond. Math. Soc. **39** (2007), no. 4, 667–676.
- [14] Y. I. Manin, *Quantum groups and non-commutative geometry*, Tech. report, Centre de Recherches Mathématiques, Université de Montreal, 1988.
- [15] A. Polishchuk and L. Positselski, *Quadratic algebras*, University Lecture Series, vol. 37, American Mathematical Society, Providence, RI, 2005.
- [16] D. R. Stephenson, *Artin-Schelter regular algebras of global dimension three*, J. Algebra **183** (1996), 55–73.
- [17] ———, *Algebras associated to elliptic curves*, Trans. Amer. Math. Soc. **349** (1997), 2317–2340.
- [18] J. Tate and M. Van den Bergh, *Homological properties of Sklyanin algebras*, Invent. Math. **124** (1996), 619–647.

□

E-mail address: benoit.kriegk@univ-st-etienne.fr

FACULT DES SCIENCES ET TECHNIQUES, LABORATOIRE DE MATHÉMATIQUES DE L'UNIVERSITÉ DE SAINT-ETIENNE, 23, RUE DU DOCTEUR PAUL MICHELON, 42023 SAINT-ETIENNE CEDEX 02

E-mail address: michel.vandenbergh@uhasselt.be

UNIVERSITEIT HASSELT, UNIVERSITAIRE CAMPUS, 3590 DIEPENBEEK